

Selective Cellular Expression of Tissue Factor in Human Tissues

Implications for Disorders of Hemostasis and Thrombosis

Thomas A. Drake,* James H. Morrissey,† and
Thomas S. Edgington†

From the Department of Pathology, UCLA, Los Angeles,*
and the Vascular Biology Group, Department of
Immunology, Research Institute of Scripps Clinic,
La Jolla,† California

Tissue factor (TF), the primary cellular initiator of the coagulation protease cascade, is implicated in having important roles in hemostasis, thrombogenesis, inflammation, and the cellular immune response, although the cytologic distribution of TF in tissues has yet to be described. This study used epitope-defined monoclonal antibodies to human tissue factor for immunohistochemical localization of TF in normal human tissues. TF was selectively expressed in tissues and was associated with cells rather than extracellular matrix. It was anatomically sequestered from blood, being undetectable in endothelium and peripheral blood cells. TF was present in vascular adventitia, organ capsules, epidermis, and mucosal epithelium. Most dermal and submucosal fibroblasts were negative. Except for alveolar macrophages and possibly dendritic cells of some lymphoid follicles, tissue macrophages did not express TF; (expression was demonstrable in LPS stimulated monocytes). Cerebral cortex, renal glomeruli, and cardiac myocytes were additional sites of prominent TF expression. Based on the cellular distribution of TF, it is hypothesized that intravascular initiation of coagulation requires induced expression by intravascular cells, and that the normal distribution of TF represents a hemostatic "envelope" ready to activate coagulation when vascular integrity is disrupted. (Am J Pathol 1989, 134:1087-1097)

Tissue factor (TF), the primary cellular initiator of the coagulation protease cascade, is a cell membrane-associated protein that serves as the receptor and essential cofactor for factors VII and VIIa.¹ The bimolecular complex of TF

and factor VII or VIIa activates factors IX and X by limited proteolysis, leading ultimately to thrombin generation and fibrin formation. As a potent initiator of coagulation, TF is believed to have a critical function in hemostasis and thrombogenesis.²⁻⁴ Cytokines, immune complexes, and other physiologic and pathologic mediators can induce TF expression in monocytes or endothelial cells, suggesting an integral involvement of TF in the pathologic processes of inflammation and thrombogenesis.⁴⁻¹³ In addition, TF is implicated in the effector phase of the cellular immune response and in the pathogenesis of certain infections.¹⁴⁻²⁰

Knowledge of the localization of TF in tissues, particularly within the vasculature, is necessary to understand the pathobiology of the thrombotic, hemorrhagic, and a variety of diseases where local activation of coagulation is indicated by the presence of fibrin. Ubiquitous tissue distribution of TF has been suggested by the constitutive production of TF by fibroblasts, smooth muscle cells, and other selected cells in culture, and the recovery of TF from various organs.^{1,21-23} Contact between TF and plasma coagulation factors does not appear likely, however, because the coagulation system is normally not activated to a significant degree. Among normal cell types, lack of expression of TF activity has been found in all circulating blood cells and endothelial cells in culture.²²⁻²⁵ Of these, only monocytes and endothelium can be induced to express TF by various mediators. Demonstration of these latter phenomena has altered the prevailing view of thrombogenesis from one of exposure of preformed TF in vascular cells with injury to one in which vascular cells play an active role by synthesizing and expressing TF and modulating other procoagulant properties in response to specific stimuli.⁴ The delineation of the *in situ* expression (or lack thereof) of TF within normal vessels and extravas-

Supported in part by NIH PO1 HL-16411, and PO1 CA-41085 and by a grant-in-aid from the American Heart Association, in part by the Florida Affiliate. This work was done during the tenure of an Established Investigatorship (JHM) from the American Heart Association.

Accepted for publication February 15, 1989.

Address reprint requests to Dr. Thomas S. Edgington, IMM-17, Research Institute of Scripps Clinic, 10666 North Torrey Pines Road, La Jolla, CA 92037.

cular tissues that is necessary to provide a morphologic foundation for any proposed mechanism of thrombogenesis and hemostasis has not been established, however.

Until recently, lack of precise delineation of human TF or reagents for its cellular localization had hampered efforts to elucidate the role of tissue expression of TF in hemostasis and in thrombotic and inflammatory diseases. Recently, TF has been purified from human brain, the cDNA cloned, and monoclonal antibodies derived to the purified protein.²⁶⁻³¹ These studies have shown that human TF is a novel protein, lacking significant homology with other known proteins. Three monoclonal antibodies that react with at least two distinct epitopes on TF, two reacting with a functional region of TF and one to a non-functional region, were used for immunohistochemical studies of TF distribution in normal human tissues. Based on the cellular distribution of TF, it is hypothesized that intravascular initiation of the coagulation protease cascade in thrombogenesis, disseminated intravascular coagulation, and related pathology requires induced expression by intravascular cells, and that TF represents a hemostatic extravascular "envelope" to which exposure of plasma proteins on disruption of vascular integrity leads to activation of coagulation. This hypothesis and the pattern of TF distribution also can account for the pattern of bleeding observed in hemophilia.

Materials and Methods

Monoclonal Antibodies

Three murine monoclonal antibodies (MAb) reactive with human TF (TF9-9C3, TF9-9B4, and TF9-10H10) were selected from a panel of MAbs derived from immunizations with purified human brain TF, the isolation and characterization of which are reported elsewhere.²⁷ Specificity for TF has been confirmed by dot blot and Western blot reactivity. Two of the antibodies recognize functional epitopes on TF, inhibiting TF initiated coagulation of human plasma by more than 90% and inhibiting specific binding of factor VII to cultured cells expressing TF by more than 80%. The third antibody recognizes a distinct nonfunctional epitope, and was used to ensure that TF would still be recognized in tissues that may have bound factor VII *in vivo*.³² All three antibodies bind strongly to cultured human cell lines expressing TF, while reactivity with cells not expressing TF is absent.

For immunohistochemical studies, the antibodies used were from hybridoma culture supernatants, diluted to working concentrations of approximately 0.1 μ g/ml in reagent diluent (2% bovine serum albumin, 0.05% thimerisol in 0.05 M TRIS-HCl, 0.9% NaCl, pH 7.4). An irrelevant IgG1K mouse MAb (ATCC TIB-115)³³ was used as a con-

trol, at equivalent concentrations in reagent diluent, also from hybridoma culture supernatant. Monoclonal antibody GCA7, specific for smooth muscle actin,³⁴ was purchased from Enzo Biochem (New York, NY).

Tissue Procurement and Preparation

Unfixed tissues were obtained from nonpathologic portions of surgical or autopsy specimens from the UCLA Medical Center. Samples were obtained within 6 hours of surgical removal (usually within 1 to 2 hours), during which time specimens were kept at 4 C, or in the case of autopsy material, within 24 hours of death. Three-millimeters thick sections were embedded in O.C.T. compound in 15 \times 15 mm cryomolds, and rapidly frozen by immersion in an isopentane-dry ice bath, then stored at -70 C. The age of patients from whom tissue was used ranged from 17 to 65 years, except for thymic tissues, which were from children under 10 years.

Immunohistochemistry

Immunohistochemistry was performed on freshly prepared cryostat sections (4 to 6 μ thickness) using a modified ABC technique.^{35,36} Sections were fixed in 2% paraformaldehyde in phosphate-buffered saline (PBS, 0.01 M sodium phosphate, 0.14 M NaCl, pH 7.4) for 2 to 5 minutes, immersed in 0.1 M glycine in 0.005 M TRIS-HCl, 0.9% NaCl (TBS wash) for 10 minutes, rinsed in distilled water for 30 seconds, and air dried for 20 to 30 minutes. Before primary antibody incubation, sections were incubated sequentially in avidin (10 U/ml) and d-biotin (1 mg/ml) for 20 minutes each (each followed by three 2-minute washes in TBS) to block endogenous avidin and biotin binding activities,³⁷ and in blocking solution (5% dry milk solids, 1% horse serum, 0.1% sodium azide in 0.01 M TRIS-HCl, 0.14 M NaCl, pH 7.4) for 20 minutes to reduce nonspecific binding. Primary antibody was incubated on sections in humidified chambers for 60 minutes at 37 C, followed sequentially by biotinylated horse antiserum IgG, H+L (Vector Laboratories, Burlingame, CA, 1:400 in reagent diluent) for 30 minutes at 37 C and streptavidin-horseradish peroxidase complex (DETEK 1-hrp, Enzo Biochemical, 1:100 in reagent diluent) 30 minutes at 37 C, with each step followed by 3 TBS washes over 10 minutes. Bound peroxidase was detected using the chromogenic substrate 3-amino-9-ethyl carbazole (0.02% in a freshly prepared solution of 0.02 M sodium acetate buffer, pH 5.1, and hydrogen peroxide, 0.03%) incubated for 10 minutes at 37 C. Sections were then washed in distilled water, counterstained with Mayer's hematoxylin, and mounted in glycerol-gelatin. Some tissues were also ex-

aminated using a commercial kit (ABC Elite, Vector Laboratories, Burlingame, CA) according to the manufacturers instructions. Sequential sections of each tissue were stained with the control MAb in parallel with specific antibody to assess nonspecific staining, which in most instances was negligible.

Peripheral blood leukocytes were prepared from buffy coat layers of freshly drawn heparinized blood. After hypotonic lysis of erythrocytes,³⁸ cells were resuspended in Medium 199 plus 10% fetal bovine serum at approximately 5×10^5 cells/ml. Cytocentrifuge preparations on poly-L-lysine-coated slides were made directly or after incubation of cell suspensions with 1 μ g/ml LPS (*Escherichia coli* 0111:B4, List Biologicals, Campbell, CA) at 37 C for 4 hours in 5% CO₂. Slides were immediately fixed in 2% paraformaldehyde plus 0.2% glutaraldehyde in PBS for 60 minutes at 4 C, permeabilized by immersion in 100% methanol at -20 C for 60 seconds, then stained as described above except for the following: slides were not air dried; 5% nonfat dry milk was included in all TBS washes; endogenous peroxidase was blocked by incubation of slides with 0.1 M periodic acid for 5 minutes then 0.02% sodium borohydride for 2 minutes, performed after the secondary antibody step.³⁹

TF Quantitation in Muscle Tissues

Skeletal and cardiac muscle were collected as described above and frozen at -70 C before use, then ground to a fine powder in a mortar and pestle in liquid nitrogen. The homogenate was diluted in 4 parts by weight of 0.05 M TRIS, 0.9% NaCl, pH 7.4 (TRIS-saline buffer) and filtered through a 100 μ steel mesh. Total protein concentrations of the filtrates were determined using a dye binding assay (Bio-Rad Laboratories, Richmond, CA).⁴⁰ Procoagulant assays were performed on samples diluted to 10 μ g of protein per ml in TRIS-saline buffer, using a plasma recalcification time technique.²² TF was quantitated by reference to a standard curve prepared using purified human brain TF reconstituted in phospholipid vesicles. These assays were performed in duplicate, with addition of 10 μ g/ml of either an inhibitory monoclonal antibody to human TF or control monoclonal to the muscle homogenates prior to assay.

Results

Specificity and General Characteristics of Staining

Evaluation of selected tissues (spleen, brain, skin, kidney, placenta) with each MAb alone and with the three in com-

Table 1. TF Procoagulant Activity in Skeletal and Cardiac Muscle

	ng TF/mg protein*	Fraction inhibited by anti-TF MAb
Cardiac muscle (N = 3)	119 \pm 65	93.9%
Skeletal muscle (N = 3)	7 \pm 4	91.5%

bination yielded identical results. This colocalization of MAbs directed against at least two distinct epitopes, including one involved in functional association with Factor VII, gave confidence that none of the three MAbs were reacting with unsuspected crossreactive epitopes on other molecules, or that the epitopes were obscured by association with Factor VII, the ligand for TF. Supportive evidence that the molecule detected immunohistochemically was functional TF was obtained by quantitating TF procoagulant activity in skeletal and cardiac muscle. Cardiac muscle, which was positive by immunohistochemistry, had 17.5 times more procoagulant activity than skeletal muscle, which was negative immunohistochemically (Table 1). Performance of the assay in the presence of a monoclonal antibody inhibiting TF function, showed that greater than 90% of the procoagulant activity measured was due to TF.

In preliminary studies, the potential effect of tissue storage at 4 C for varying periods of time before freezing was examined. Portions of skin, brain, and placenta received within 60 minutes of removal from the body were frozen immediately and after 8 and 24 hours while being kept moist at 4 C. Immunohistochemistry performed at one time on sections from each showed no evidence of either loss or new appearance of antigen over time. Also, no differences were noted in the patterns of reactivity or relative intensity between like tissues from surgical or autopsy sources. These observations indicated that the TF epitopes detected were relatively stable in tissues within the range of conditions encountered in this study.

Immunohistochemical localization of TF showed differential expression among the various tissues examined (Table 2, Figures 1-17). No substantial differences were noted between tissues from different individuals. A general finding was that TF appeared to be cell associated, within the resolution of light microscopy. At no site was there evidence for extracellular matrix localization. Chromogen localization in positive cells was usually diffuse rather than focal or limited to the cell outline.

Blood Vessels

TF in blood vessels was predominately localized to cells of the adventitia, being undetectable in endothelium and

Table 2. Summary of TF Expression in Human Tissues

Skin	Epidermis	+++
	Dermis	—
Gut	Mucosa	+++
	Submucosa	—
	Muscularis	V (+)
Vessels	Intima	—
	Media	V (+)
	Adventitia	++
	Capillaries	—
Heart	Myocardium	+++
	Endocardium	—
	Cardiac valves	—
Lung	Bronchial mucosa	++
	Bronchial submucosa	—
	Alveolar septae	+
	Alveolar epithelial cells	++
	Alveolar macrophages	V (++)
Brain	Meninges	+
	Cerebral cortex	+++
Kidney	Glomeruli	+++
	Tubules	—
	Interstitial	—
Spleen	Capsule	++
	Trabeculae	+++
	Splenic cords	—
	Lymphoid areas	—
Liver	Hepatocytes	+
	Kupffer cells	—
	Biliary duct epithelium	—
Adrenal glands	Cortex	—
	Medulla	+
Peripheral nerve	Schwann cells	++
	Axons	—
Skeletal muscle	Myocytes	—
	Perimysium	—

—, absent; +, weak intensity; ++, moderate intensity; +++, strong intensity; V, variably present, with intensity, when present, in parentheses.

variably present in cells composing the media (Figures 6,7,9,18,19). Aorta and elastic arteries had prominent adventitial localization that was distinctly cell associated and not present in connective tissue matrix. The medial layer of these vessels showed variable weak to moderate reactivity in some areas. Intima was consistently negative. Most muscular arteries and larger arterioles also had distinct adventitial reactivity, although in some locations (kidney and gut submucosa) this was absent or minimal but in other locations (spleen, superficial cerebral vessels) was

exceptionally intense. (These differences were consistent among specimens from different sources and could not be attributed to individual variation or to sample handling.) Cells in the media of these vessels and in smaller arterioles was generally negative, although they were occasionally weakly to moderately positive, in focal areas. Smaller arterioles also had adventitial cell localization but this was often not prominent and, as with larger vessels, was absent in kidney and gut.

Capillaries and postcapillary venules did not react at any site. Adventitial cells of small to medium sized veins generally reacted similarly to those of arteries of comparable size and location. In large veins, TF was expressed in cells that were interspersed between bundles of smooth muscle cells. Because a defined medial layer is thin to nonexistent in these vessels,⁴¹ TF expressing cells were irregularly present near the intima.









Parallel sections of larger arteries and veins were reacted with MAb to TF and to smooth muscle actin to determine whether the TF positive cells in adventitia were actually of smooth muscle origin. In both arteries and veins, adventitial cells expressing TF did not contain immunoreactive smooth muscle actin, suggesting they were fibroblasts (Figures 18, 19).

An exception to the absent or irregular reactivity of vascular smooth muscle was observed in umbilical cord and placental arteries and veins, where medial smooth muscle cells expressed TF uniformly with moderate intensity. Endothelium was negative, whereas adventitial cells reacted intensely, as did the stromal cells in the umbilical cord matrix.

Peripheral Blood Cells

Unstimulated peripheral blood leukocytes were uniformly negative (Figure 8a). Incubation of cells with endotoxin for 4 hours induced immunohistochemically detectable TF in monocytes but not neutrophils or lymphocytes (Figure 8b). Eosinophils, basophils, erythrocytes, and platelets also were negative. Lymphocytes in spleen, thymus, and lymph nodes were negative. Macrophages in tissues and derived cells lining vascular channels in spleen and liver

Figure 1. Skin (inset: control MAb): epidermis is strongly positive while dermis is negative (X231). **Figure 2.** Colonic mucosa (inset: control MAb): epithelium is positive while lamina propria is negative except for cells of the subepithelial fibroblastic sheath (X231). **Figure 3.** Vaginal mucosa: association of TF expression with maturation of the squamous epithelium is marked (X92). **Figure 4.** Bronchial mucosa (small bronchus): epithelium is positive, staining more strongly in basally located cells; adjacent submucosa is negative (X369). **Figure 5.** Urinary bladder mucosa: intense staining occurs in upper submucosa whereas transitional epithelium is only faintly positive (X231). **Figure 6.** Large meningeal artery (L, lumen): adventitial cells are strongly positive while media and intima are negative (X231). **Figure 7a.** Carotid artery (L, lumen): adventitia is positive, media weakly positive, and intima negative (X92). **b:** Iliac vein (L, lumen): scattered cells in the adventitia are positive, the intima is negative (X231). **Figure 8a.** Peripheral blood leukocytes (unstimulated): neutrophils, lymphocytes and monocytes (M) are all negative (X577). **b:** Peripheral blood leukocytes exposed to LPS for 4 hours: a monocyte (M) has been induced to express TF whereas neutrophils and lymphocytes remain negative (X577). **Figure 9a.** Arteriole, uterus: adventitia of vessel is positive; vascular intima and media and uterine smooth muscle is negative (X231). **b:** Arteriole and small vein, muscular wall of colon: no staining of adventitia is seen, in contrast with vessel in panel a; autonomic ganglion (arrow) is positive, smooth muscle of muscularis very weakly positive (X231).



also were negative. Exceptions were alveolar macrophages, which were variably positive with weak to moderate intensity, and cells of dendritic morphology in germinal centers of reactive follicles in spleen, tonsil, and gut.

Extravascular Tissues

TF was selectively expressed in various organs and tissues as summarized in Table 2. Prominent expression occurred in epidermis, bowel and respiratory mucosa, cerebral cortex, myocardium, and renal glomeruli. Also, cells composing the fibrous capsules of organs such as liver, spleen, kidney, and adrenals were consistently positive (Figure 13). Expression of TF by connective tissue fibroblasts was relatively infrequent, occurring most prominently in the adventitia of blood vessels, and in the subepithelial fibroblastic sheath of bowel mucosa. Fibroblasts in connective tissues elsewhere, whether densely collagenous, as in dermis or joint capsules (Figure 11), or loose, as in submucosal tissues and the perimysium of skeletal and cardiac muscle, were generally negative.

Epidermis reacted intensely, with the strongest expression consistently observed in the granular cell layer, whereas the basal layer was weak to negative (Figure 1). This pattern also was observed in the squamous epithelium of the oropharynx and vagina (Figure 3). Dermal connective tissue and capillaries did not react. Respiratory, intestinal, and urinary bladder mucosae were positive, although the intensity of the reaction varied. Mucosal epithelium and derived submucosal glands of larger bronchi were moderately to strongly positive, whereas epithelial reactivity in smaller airways was more variable (Figure 4). In the bowel, epithelium reacted with weak to moderate intensity (Figure 2). Throughout the bowel, however, the most intense localization was present in the 1 to 2 cell thick layer of the subepithelial fibroblastic sheath (pericryptal fibroblasts) located immediately subjacent to the epithelium. Other cells in the lamina propria did not express TF. In the urinary bladder, transitional epithelium was weakly positive and in some samples, negative (Figure 5). Intense reactivity occurred in a band of otherwise nondescript upper submucosal stromal cells, whereas cells deeper in the submucosa were negative.









TF was strongly expressed in cerebral cortex, being present diffusely throughout (Figure 12). Neuronal and other cell bodies were positive, but not more prominently than the surrounding neuropil. Cardiac myocytes expressed TF with a uniform cytoplasmic distribution (Figure 10a). This was in striking contrast with skeletal muscle from various sites, which was consistently negative (Figure 10b), as was smooth muscle in most locations. In renal parenchymal structures, TF was limited to the glomeruli (Figure 15). Epithelial and mesangial cells of the glomerular tuft reacted strongly, whereas glomerular capillary endothelium was negative. Epithelium of Bowman's capsule also was positive.

Discussion

Maintenance of a potent, rapidly active hemostatic system is a physiologic necessity that must be structured in a manner that minimizes the risk of intravascular coagulation. This is achieved by limiting activation of coagulation, inhibiting the activated proteases and cofactors that are formed, and by active fibrinolysis.⁴² Pathologic activation of coagulation is key to thrombotic diseases and in inflammation. TF is a major cellular initiator of the coagulation protease cascade, functioning as the cell surface receptor and essential cofactor for Factor VII.¹ Because contact of TF with blood is sufficient to initiate coagulation, TF expression in vessels must be strictly regulated to preclude activation. This study shows that TF is anatomically sequestered from the blood in the normal state, being selectively expressed at sites isolated from plasma. Normal peripheral blood cells and endothelium did not have detectable TF, nor did smooth muscle cells of most vessels. TF associated with vessels was predominantly localized to cells in the adventitia.

The absence of detectable TF in endothelium and other cells in contact with blood is consistent with *in vitro* and *ex vivo* studies, which have shown negligible activity in unstimulated endothelium or leukocytes.²²⁻²⁵ TF-initiated thrombosis is therefore unlikely to result from simple injury to normal endothelium with loss of cellular integrity, in the absence of previous "activation." Endogenous inflammatory mediators such as interleukin-1 and tumor necrosis factor, as well as endotoxin and immune com-

Figure 10a. Cardiac muscle: myocytes are positive (X231). b. Skeletal muscle: myocytes are negative (X231). Figure 11. Synovium, knee: synovial cells and connective tissues of stroma are negative or only weakly positive in vicinity of small vessels (X231). Figure 12. Cerebral cortex: cortical parenchyma is strongly positive in diffuse pattern; meninges are moderately positive; adventitial staining is present around a small meningeal vessel (X92). Figure 13a. Lymph node, cortex: capsule is moderately positive, lymphocytes are negative (X369). b. Liver: capsule is moderately positive, whereas hepatocytes are weakly positive and Kupffer cells negative (X369). Figure 14. Lung: alveolar macrophages and epithelial cells are positive (X369). Figure 15. Glomerulus, kidney: glomerular interstitium and cells of Bowman's capsule are positive; endothelium of capillary loops appear negative, as do surrounding renal tubules (X231). Figure 16. Placental villi: trophoblast is negative whereas internal stroma is positive, strongest around vessels (X231). Figure 17. Spleen with arteriole and adjacent lymphoid follicle: adventitia is strongly positive as are scattered cells in the center of the follicle (X92).






Figure 18. Saphenous vein: parallel sections incubated with anti-smooth muscle actin MAb (a) and anti-TF MAb (b) showing lack of smooth muscle actin expression in TF positive cells which surround bundles of smooth muscle (I, intima; $\times 369$). **Figure 19.** Aorta, junction of media and adventitia: parallel sections incubated with anti-smooth muscle actin MAb (a) and anti-TF MAb, (b) showing lack of smooth muscle actin expression in TF positive cells in adventitia (M, media; A, adventitia; $\times 231$).

plexes, can induce endothelial cell TF expression and are implicated in the thrombogenic process, although endothelial cell expression of TF *in vivo* remains to be demonstrated.⁶⁻¹⁰ Monocytes also are capable of being induced to express TF by various stimuli and also are implicated in thrombogenesis as well as extravascular fibrin deposition in inflammation.^{5,11,13,14} Monocytes and macrophages at sites in contact with blood such as liver and spleen did not express detectable TF, nor did these cells at many extravascular sites, lung being a prominent exception.

We were able to demonstrate immunohistochemically, as has been indicated by other techniques,^{24,25} that the monocyte, when stimulated with endotoxin, an etiologic agent in septic shock, is the only normal peripheral blood cell to express tissue factor. This is consistent with previous implications of this cell as a potentially responsible initiating cell in thrombogenesis and in cellular immune responses.

The observed distribution of TF is consistent with its presumed primary function in hemostasis. TF was generally localized in what may be thought of as an "envelope" distribution: it surrounded blood vessels in adventitia, encased organs in fibrous capsules, and was present in epidermis and mucosal epithelium and subjacent stromal cells in skin and internal mucosae, thus providing a contin-

uous hemostatic barrier with the external environment. In brain, where even minor hemorrhage may be catastrophic, TF was strongly expressed throughout the cortical parenchyma.

Although there are as yet no documented cases of TF deficiency (a complete absence of which may be fatal in utero) to confirm a primary role for TF in hemostasis, collective information viewed in light of its anatomic distribution supports such a role. Although Factor VII deficiency has been known to be associated with impaired hemostasis, the defects associated with Factors VIII and IX deficiencies have implicated the contact activation pathway as being of major importance. Recent studies have shown that TF-VIIa complex can activate Factor IX as well as Factor X, and that the former pathway is responsible for the majority of TF initiated Xa formation at low TF concentrations.^{43,44} Studies by Broze et al⁴⁵ suggest that the recently described lipoprotein-associated coagulation inhibitor is responsible for the latter phenomenon. It is particularly interesting that the sites most commonly affected by spontaneous hemorrhage in Factor VIII or IX deficient patients, deep skeletal muscle and joints, had undetectable to low TF expression. Clinical studies in hemophilic patients who are unresponsive to replacement therapy because of inhibitory antibodies have shown that

infusions of Factor VIIa can stop bleeding.^{46,47} Such responses with enhancement of TF-dependent activation of coagulation suggest an important hemostatic role for the presumed low levels of TF present at those sites. Also, our observations of high levels of TF expression in epidermis are consistent with a recent study that showed activation of coagulation to be the earliest hemostatic event in bleeding time wounds.³ Such activation occurred in healthy individuals as well as in Factor VIII and IX deficient people, but was impaired in those with Factor VII deficiency, implicating TF rather than the contact activation pathway as the initiating mechanism. In this setting, the high level of TF expression in epidermis would be expected to preferentially promote direct X activation, making the response independent of IX activation, as was actually observed. Early thrombin generation may be an important mechanism of platelet activation in the primary hemostatic response in such wounds, which has until now been considered to be largely independent of the coagulation protease cascade.

Although much work remains to establish the relative importance of the TF and contact activation pathways in hemostasis, this study is fully consistent with and further supports previous observations that TF has a major role in this essential function. TF did not appear to be present in extracellular matrix, the putative site of contact activation, but rather was cell associated, thus providing the body with complementary rather than duplicated sites for coagulation activation.

The foregoing discussion assumes that the TF detected immunohistochemically in this study was functionally active. Two of the three MAbs used reacted with functional epitopes on TF. Also, tissues known to be rich in TF, such as brain and placenta, reacted strongly and we were able to demonstrate substantially more procoagulant activity, attributable to TF, in an immunohistochemically positive tissue, cardiac muscle, than in a comparable tissue that was negative, skeletal muscle. One study reported finding very little procoagulant activity in isolated renal glomeruli of normal kidneys, however, a site we consistently found to be strongly positive.⁴⁸ It is possible that the TF in glomeruli is predominately extracellular, and lacking in function because it would not be associated with membrane lipid required for activity. Further work will be necessary to reconcile these observations, and to determine whether TF at other sites of localization is functionally active.

Of importance also is the unknown sensitivity of immunohistochemistry in detecting low levels of TF in cells. This is complicated by a lack of knowledge concerning the levels of TF expression that may be physiologically significant. Studies by Dvorak and colleagues with an animal model have demonstrated extravascular fibrin deposition in dermis occurring after induction of increased vascular

permeability.⁴⁹ In conjunction with knowledge that cultured fibroblasts express tissue factor, it was postulated that TF is normally expressed by dermal fibroblasts *in vivo*. Our inability to detect TF in human dermis could reflect some limitations of sensitivity of the analyses, among other explanations that include species differences in local TF expression. Interestingly, cultured human fibroblasts express substantial levels of TF which are detectable immunohistochemically (data not shown).

The nonuniform distribution of TF throughout the body shows that TF expression is significantly regulated in various cell types. The findings in two tissues in particular were indicative of an important association of TF expression with cellular differentiation. Among muscular tissues, TF was strongly expressed in cardiac muscle, variably expressed in smooth muscle, and was undetectable in skeletal muscle. In epidermis, TF expression was absent to weak in the germinative basal layer and strongest in the more differentiated outer layers. This potential link between squamous differentiation and TF expression also was suggested by the presence of TF in Hassell's corpuscles in the absence of expression in other thymic epithelium. An important extension of the finding of differential expression of TF in normal tissues will be evaluation of its expression in their neoplastic counterparts, and the association of thrombosis and malignancy.

TF expression by cells not normally positive, either through induction by biologic mediators or loss of regulation, may be important in various biologic and pathologic processes. As discussed above, this is probably of particular importance in thrombosis and inflammation in which endothelium and monocytes can be induced to express TF. The variable expression of TF by smooth muscle and fibroblasts at different locations suggests that these cells also may modulate TF expression in response to biologic stimuli. Although much of the distribution of TF in normal tissues can be understood in terms of its procoagulant function, its selective expression at sites not of obvious importance for hemostasis raises the question as to whether it may have additional biologic functions.

References

1. Nemerson Y, Bach R: Tissue factor revisited. *Prog Hemostasis Thromb* 1982, 6:237-261
2. Nemerson Y: Tissue factor and hemostasis. *Blood* 1988, 71: 1-8
3. Weiss HJ, Lages B: Evidence for tissue factor-dependent activation of the classic extrinsic coagulation mechanism in blood obtained from bleeding time wounds. *Blood* 1988, 71: 629-635
4. Rodgers GM: Hemostatic properties of normal and perturbed vascular cells. *FASEB J* 1988, 1:116-123

5. Conkling PR, Greenberg CS, Weinberg JB: Tumor necrosis factor induces tissue factor-like activity in human leukemia cell line U939 and peripheral blood monocytes. *Blood* 1988, 72:128-133
6. Colucci M, Balconi G, Lorenzet R, Pietra A, Locati D, Donati MB, Semerara N: Cultured human endothelial cells generate tissue factor in response to endotoxin. *J Clin Invest* 1983, 71:1893-1896
7. Bevilacqua MP, Pober JS, Majeau GR, Cotran RS, Gimbrone MA Jr: Interleukin 1 induces biosynthesis and cell surface expression of procoagulant activity in human vascular endothelial cells. *J Exp Med* 1984, 160:618-623
8. Bevilacqua MP, Pober JS, Manjeau GR, Fiers W, Cotran RS, Gimbrone MA: Recombinant tumor necrosis factor induces procoagulant activity in cultured human vascular endothelium: Characterization and comparison with the actions of interleukin 1. *Proc Natl Acad Sci USA* 1986, 83:4533-4537
10. Tannenbaum SH, Finko R, Cines DB: Antibody and immune complexes induce tissue factor production in human endothelial cells. *J Immunol* 1986, 137:1532-1537
11. Prydz H: Synthesis and surface availability of thromboplastin and plasminogen activator in mononuclear phagocytes, *Mononuclear Phagocytes: Physiology and Pathology*. Edited by RT Dean, W Jessup. New York, Elsevier, 1985, pp 115-137
12. Kaplan KL: Coagulation proteins in thrombosis, Hemostasis and Thrombosis, *Basic Principles and Clinical Practice*. 2nd ed. Edited by RW Coleman, J Hirsch, VJ Marder, EW Slazman. Philadelphia, JB Lippincott, 1987, pp 1098-1110
13. Cybulsky MI, Chan MKW, Movat HZ: Acute inflammation and microthrombosis induced by endotoxin, interleukin-1, and tumor necrosis factor and their implications in gram-negative infection. *Lab Invest* 1988, 58:365-378
14. Hopper KE, Geczy CL, Davies WA: A mechanism of migration inhibition in delayed-type hypersensitivity reactions: I. Fibrin deposition on the surface of elicited peritoneal macrophages in vivo. *J Immunol* 1981, 126:1052
15. Helin H, Edgington TS: Allogenic induction of the human T cell instructed monocyte procoagulant response is rapid and is elicited by HLA-DR. *J Exp Med* 1983, 158:962-975
16. Gregory SA, Edgington TS: Tissue factor induction in human monocytes: Two distinct mechanisms displayed by different alloantigen responsive T cell clones. *J Clin Invest* 1985, 76:2440-2445
17. Geczy CL, Meyer PA: Leukocyte procoagulant activity in man: An in vitro correlate of delayed-type hypersensitivity. *J Immunol* 1982, 128:331-336
18. Levy GA, Leibowitz JL, Edgington TS: Induction of monocyte procoagulant activity by murine hepatitis virus type 3 parallels disease susceptibility in mice. *J Exp Med* 1971, 154:1150-1163
19. Ryan J, Geczy C: Coagulation and the expression of cell-mediated immunity. *Immunol Cell Biology* 1987, 127-139
20. Drake TA, Rodgers GM, Sande MA: Tissue factor is a major stimulus for vegetation formation in enterococcal endocarditis in rabbits. *J Clin Invest* 1984, 73:1750-1753
21. Kadish JL, Wenc KM, Dvorak HF: Tissue factor activity of normal and neoplastic cells: Quantitation and species specificity. *J Natl Cancer Inst* 1983, 70:551-557
22. Maynard JR, Dreyer BE, Stemerman MB, Pittick FA: Tissue factor coagulant activity of cultured human endothelial and smooth muscle cells and fibroblasts. *Blood* 1977, 50:387-396
23. Rodgers GM, Greenberg CS, Shuman MA: Characterization of the effects of cultured vascular cells on the activation of blood coagulation. *Blood* 1983, 61:1155-1162
24. Rivers RPA, Hathaway WE, Weston WL: The endotoxin-induced procoagulant activity of human monocytes. *Br J Haematol* 1975, 30:311-316
25. Edwards RL, Rickles FR, Bobrove AM: Mononuclear cell tissue factor: Cell of origin and requirements for activation. *Blood* 1979, 54:359-370
26. Morrissey JH, Revak D, Tejada P, Fair DS, Edgington TS: Resolution of monomeric and heterodimeric forms of tissue factor, the high-affinity cellular receptor for factor VII. *Thromb Res* 1988, 50:481-493
27. Morrissey JM, Fair DS, Edgington TS: Monoclonal antibody analysis of purified and cell-associated tissue factor. *Thromb Res* 1988, 52:247-261
28. Morrissey JH, Fakhrai H, Edgington TS: Molecular cloning of the cDNA for tissue factor, the cellular receptor for the initiation of the coagulation protease cascade. *Cell* 1987, 50:129-135
29. Spicer EK, Horton R, Bloem L, Bach R, Williams KR, Guha A, Kraus J, Lin T-C, Nemerson Y, Konigsberg WH: Isolation of cDNA clones coding for human tissue factor: Primary structure of the protein and cDNA. *Proc Natl Acad Sci USA* 1987, 84:5148-5152
30. Scarpati EM, Wen D, Broze GJ Jr, Miletich JP, Flandermeyer RR, Siegel NR, Sadler JE: Human tissue factor: cDNA sequence and chromosome localization of the gene. *Biochemistry* 1987, 26:5234-5238
31. Fisher KL, Gorman CM, Vehar GA, O'Brien DP, Lawn RM: Cloning and expression of human tissue factor cDNA. *Thromb Res* 1987, 48:89-99
32. Ruf W, Morrissey JM, Edgington TS: Monoclonal antibodies inhibit tissue factor function by two distinct mechanisms: Evidence for a functional region of tissue factor not involved in factor VII binding. (Submitted for publication)
33. Gurney EG, Harrison RO, Fenno J: Monoclonal antibodies against simian virus 40 T antigens: Evidence for distinct subclasses of large T antigen and for similarities among nonviral T antigens. *J Virol* 1980, 34:752-763
34. Gown AM, Vogel AM, Gordon D, Lu PL: A smooth muscle-specific monoclonal antibody recognizes smooth muscle actin isozymes. *J Cell Biol* 1985, 100:807-813
35. Guesdon JL, Ternynck T, Avrameas S: The use of avidin-biotin interaction in immunoenzymatic techniques. *J Histochem Cytochem* 1979, 27:1131-1139
36. Hsu S-M, Raine L, Fanger H: The use of avidin-biotin-peroxidase complex (ABC) in immunoperoxidase techniques: A comparison between ABC and unlabeled antibody PAP procedures. *J Histochem Cytochem* 1981, 29:577-580
37. Wood GS, Warnke R: Suppression of endogenous avidin-binding activity in tissues and its relevance to biotin-avidin detection systems. *J Histochem Cytochem* 1981, 29:1196-1204

38. Metcalf JA, Gallin JI, Nauseef WM, Root RK: Laboratory Manual of Neutrophil Function. New York, Raven Press, 1986, p 3
39. Heyerman E, Neville AM: A shorter immunoperoxidase technique for demonstration of carcinoembryonic antigen and other cell products. *J Clin Pathol* 1977, 30:138-140
40. Bradford M: A rapid and sensitive method for quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Ann Biochem* 1976, 72:248-254
41. Rhodin, JAG: Architecture of the vessel wall, Handbook of Physiology, Section 2, The Cardiovascular System. Volume II: Vascular Smooth Muscle. Edited by DE Bohr, AP Somlyo, HV Sparks Jr. Bethesda, American Physiological Society, 1980, pp 1-31
42. Coleman RW, Marder VJ, Salzman EW, Hirsh J: Overview of hemostasis, Hemostasis and Thrombosis, Basic Principles and Clinical Practice. 2nd ed. Edited by RW Coleman, J Hirsch, VJ Marder, EW Salzman. Philadelphia, JB Lippincott, 1987, pp 3-17
43. Osterud B, Rapaport S: Activation of factor IX by the reaction product of tissue factor and factor VII: Additional pathway for initiating blood coagulation. *Proc Natl Acad Sci USA* 1977, 74:5260-5264
44. Marler RA, Kliess AJ, Griffin JH: An alternative extrinsic pathway of human blood coagulation. *Blood* 1982, 60:1353-1358
45. Broze GJ Jr, Warren LA, Novotny WF, Higuchi DA, Girard JJ, Militich JP: The lipoprotein-associated coagulation inhibitor that inhibits the factor VII-tissue factor complex also inhibits factor Xa: Insight into its possible mechanism of action. *Blood* 1988, 71:335-343
46. Lusher JM, Shapiro SS, Palascak JE, Rao AV, Levine PH, Blatt PM, Hemophilia Study Group: Efficacy of prothrombin complex concentrates in hemophiliacs with antibodies to factor VIII. *N Engl J Med* 1980, 303:421-425
47. Hedner U, Kisiel W: Use of human factor VIIa in the treatment of two hemophilia A patients with high-titer inhibitors. *J Clin Invest* 1983, 71:1836-1841
48. Tipping PG, Dowling JP, Holdsworth SR: Glomerular procoagulant activity in human proliferative glomerulonephritis. *J Clin Invest* 1988, 81:119-125
49. Dvorak HF, Senger DR, Dvorak AM, Harvey VS, McDonagh J: Regulation of extravascular coagulation by microvascular permeability. *Science* 1985, 227:1059-1061